

DISORIENTED CHIRAL CONDENSATE AND CHARGE-NEUTRAL
PARTICLE FLUCTUATIONS IN HEAVY ION COLLISIONS

MLADEN MARTINIS,¹ and VESNA MIKUTA-MARTINIS²

*Theoretical Physics Division
Rudjer Bošković Institute, P.O.B. 1016,
41001 Zagreb, CROATIA*

Abstract

The possibility of large charge and isospin fluctuations in high-energy heavy ion-collisions is studied within the framework of the nonlinear σ -model with quark degrees of freedom. The multipion exchange potential between two quarks is derived. It is shown that the soft chiral pion bremsstrahlung leads to anomalously large fluctuations in the ratio of neutral to charged pions, predicted by the formation of a disoriented chiral condensate (DCC). The factorization property of the scattering amplitude in the impact parameter space of the leading two-nucleon system is used to study semiclassical production of pions in the central region. We show that the DCC-type fluctuations are suppressed if a large number of pions are produced in ρ -type clusters. Our conclusion is supported through the calculation of two pion correlation parameters as a function of the ρ -to- π ratio.

PACS numbers: 25.75.+r, 12.38.Mh, 13.85.Tp, 24.60.Ky

¹e-mail address: martinis@rudjer.irb.hr

²e-mail address: vmikuta@rudjer.irb.hr

1. *Introduction*

Central ultrarelativistic collisions at RHIC and LHC with more than 3000 produced particles present remarkable opportunity to analyse event-by-event fluctuations of hadronic observables. Such single events analysis with large statistics can reveal new physical phenomena usually hidden when averages over a large statistical sample of events are made [1,2]. The numbers of particles produced in relativistic heavy-ion collisions can differ dramatically from collision to collision due to the variation of impact parameter (centrality dependence), energy deposition (leading particle effect), and other dynamical effects [2,3]. The fluctuations can also be influenced by novel phenomena such as disoriented chiral condensate (DCC) [4,5].

The best probes of such novel dynamics are fluctuations of conserved quantities, because conservation laws limit the degree to which final-state scattering can dissipate. Even globally conserved quantities such as energy, net charge, isospin, baryon number and strangeness can fluctuate when measured, e.g., in a limited phase space region.

Several methods have been proposed [3,7] to distinguish between statistical and dynamic fluctuations. In high energy hadronic and heavy-ion collisions the correlations of particle production are usually analysed and described in terms of "short range correlations" (SRC) and "long range correlations" (LRC). The SRC are mainly due to clustering of outgoing particles. These clusters may be hadronic resonances which decay into a few particles or minijets [5,6,8]. The LRC dominate in high multiplicity events and are mainly due to global conservation laws.

The old puzzle in cosmic-ray observations is the existence of few exotic events characterized by an anomalously large number of charged pions in comparison with the number of neutral pions, the Centauros [9], indicating that there should exist a

strong negative long-range correlation between two types of the pions.

Such long-range correlations are possible if pions are produced semiclassically and constrained by global conservation of isospin [10–15].

Although the actual dynamical mechanism of the production of a classical pion field in the course of a high-energy collision is not known, there exist numerous interesting recent theoretical attempts to explain Centauros either as different types of isospin fluctuations due to the formation of a disoriented chiral condensate (DCC) [16–20], or as multiparticle Bose-Einstein correlations (BEC) [21], or as the formation of a strange quark matter (SQM) [22]. Among the most interesting speculations is the idea of DCC that localized regions of misaligned chiral vacuum might occur during the ultrahigh-energy hadronic and heavy-ion collisions when the chiral symmetry is restored at high temperatures. These regions, if produced, would behave as a pion laser, relaxing to the ground state by coherent pion emission []. It is generally accepted that the fluctuation of the ratio of neutral to charged pions of the Centauro type could be a sign of the DCC formation provided that a single large domain is formed, containing a large number of low p_T pions. Since the pions formed in the DCC are essentially classical they form a quantum superposition of coherent states with different orientation in isospin space. If all the pions in the domain are pointing in the same isospin direction and the condensate state is a pure isoscalar then the formation of DCC leads to large event-by-event fluctuations in the ratio $f = n_0/n$ of the number of π_0 's in the DCC divided by the total number of pions produced in an event. The probability distribution of f inside the DCC domain is [13,16,17,19]

$$P_{DCC}(f) = \frac{1}{2\sqrt{f}} \quad (1)$$

There are a variety of proposed mechanisms other than DCC which also lead to the distribution (1) [23,24,25,26]. The distribution $P_{DCC}(f)$ is different from the generic

binomial-distribution expected in normal events which assumes equal probability for production of π_+ , π_- and π_0 pions. The emission of charged and neutral pions is then uncorrelated and the distribution

$$P_B(n_0, n) = \binom{n}{n_0} \left(\frac{1}{3}\right)^{n_0} \left(\frac{2}{3}\right)^{n-n_0} \quad (2)$$

in the limit as $n \rightarrow \infty, n_0 \rightarrow \infty$ with f fixed, approaches a delta function at $f = 1/3$.

The possibility of observing the DCC type fluctuations critically depends on the size and the energy content of the DCC domain. If the domain is of the pion size, the effect of DCC is too small to be observed experimentally. The early accelerator searches for Centauros and DCC at CERN [27–30] and at Fermilab [23,31] were thus unsuccessful. With the RHIC facility at BNL now, there is a possibility to consider event-by-event fluctuations of other specific hadronic observables that might be more informative about the signal of DCC. There are various factors that may affect the observation of the DCC signal [32]. Experimental signals such as the isospin fluctuations, the strong relative enhancement of the number of low p_T pions, and the suppression of HBS correlations may provide a robust signal of DCC [33].

The space-time scenario of the formation and decay of the DCC is usually studied within one of the simplified versions of the chiral effective Lagrangians, either the linear or nonlinear sigma model [20]. However, it should be emphasized that the use of σ -models, be they linear or nonlinear, is only a rough approximation to the true dynamics, because the couplings of pion and sigma fields to the constituent quarks may be large and their effect should not be ignored.

In this paper, we present results of an event-by-event analysis of charge-neutral pion fluctuations as a function of the ρ/π ratio in pp-collisions. Following the approach of our earlier papers [24,34], we consider in Section 2. the leading-particle

effect as a source of a classical pion field in the impact parameter space which is related to the semiclassical solutions of the equation of motion of the nonlinear σ -model coupled to quark degrees of freedom. In order to facilitate the analysis of charged-neutral pion correlations, we also derive the corresponding pion-generating function. The quantum equations of the nonlinear σ -model coupled to quarks are discussed in Section 3., and shown to lead to a coherent state description of the pions emerging from the DCC. A multipion exchange potential between two quarks is derived as well as a corresponding correlator for multipion production. Results of our investigation are summarized in the Section 4. Our general conclusion is that within the nonlinear σ model the large isospin fluctuations depend strongly on the value of the ρ/π ratio which fluctuate from event to event.

2. Pion production from a classical source

At high energies most of the pions are produced in the nearly baryon-free central region. The energy available for the hadron production is

$$E_{had} = \frac{1}{2}\sqrt{s} - E_{leading} \quad (3)$$

which at fixed total c.m. energy $\sqrt{s}$ varies from event to event. The n-pion contribution to the s -channel unitarity can be written as an integral over the relative impact parameter b of the two incident leading particles:

$$F_n(s) = \frac{1}{4s} \int d^2b \prod_{i=1}^n dq_i |T_n(s, \vec{b}; 1 \dots n)|^2, \quad (4)$$

where $dq = d^2q_T dy / 2(2\pi)^3$. The normalization is such that

$$\begin{aligned} F_n(s) &= s\sigma_n(s), \\ \sigma_{inel}(s) &= \sum_{n=1}^{\infty} \sigma_n(s). \end{aligned} \quad (5)$$

If the isospin of the incoming leading particles is II_3 , then the initial-state vector of the pion field is $\hat{S}(s, \vec{b}) |II_3\rangle$, where $|II_3\rangle$ denotes a vacuum state with no pions.

The n -pion production amplitude is

$$iT_n(s, \vec{b}; q_1 \dots q_n) = 2s \langle I' I'_3; q_1 \dots q_n | \hat{S}(s, \vec{b}) | II_3 \rangle, \quad (6)$$

where $I' I'_3$ denotes isostate of the outgoing leading particles.

The basic assumption of the independent emission of pion-clusters, in b -space is the factorization property of the scattering amplitude. The factorization of T_n follows if the quantum field of the pion-cluster satisfies the equation of motion of the form

$$(\square + m_c^2) \vec{\pi}_c(s, \vec{b}; x) = \vec{j}_c(s, \vec{b}; x), \quad (7)$$

for an isovector cluster and similar one for an isoscalar cluster. Here, $\vec{j}_c$ is a classical source of the cluster which decay into $c = 1, 2, \dots$ pions outside the region of strong interactions (the final-state interaction between pions being neglected). Clusters decaying into two or more pions simulate a short-range correlation between pions. They need not be well-defined resonances.

If the conservation of isospin is a global property of the colliding system, then $\vec{j}_c(s, \vec{b}; x)$ is of the form

$$\vec{j}_c(s, \vec{b}; x) = j_c(s, \vec{b}; x) \vec{n}, \quad (8)$$

where $\vec{n}$ is a fixed unit vector in isospace independent of x, s and $\vec{b}$. The global conservation of isospin thus introduces the long-range correlation between the emitted pions.

The S matrix following from such a classical source is still an operator in the space of pions. Inclusion of isospin requires $\hat{S}(s, \vec{b})$ to be also a matrix in the isospace of the leading particles.

The coherent production of pion-clusters is described by the following S matrix:

$$\hat{S}(s, \vec{b}) = \int d^2 \vec{n} | \vec{n} \rangle \hat{D}(s, \vec{b}) \langle \vec{n} |, \quad (9)$$

where $|\vec{n}\rangle$ represents the isospin-state vector of the two-leading-particle system. The quantity $\hat{D}(s, \vec{b})$ is the unitary coherent-state displacement operator defined in our case as

$$D(s, \vec{b}) = \exp[a^\dagger(s, \vec{b}) - a(s, \vec{b})] \quad (10)$$

where

$$a^\dagger(s, \vec{b}) = \sum_c \int dq [J_c(s, \vec{b}; q) \vec{n} \vec{a}_c^\dagger(q) + J'_c(s, \vec{b}; q) a_c^\dagger(q)], \quad (11)$$

and $\vec{a}_c^\dagger(q)$ and $a_c^\dagger(q)$ are the creation operators of a cluster of type c , respectively.

The isospin (I', I'_3) of the outgoing leading particle system varies from event to event. It is produced with the probability ω_{I', I'_3} , and we can sum over all (I', I'_3) to obtain the probability distribution of producing $n_+ \pi^+$, $n_- \pi^-$, and $n_0 \pi^0$ from a given isospin state:

$$P_{II_3}(n_+ n_- n_0) N_{II_3} = \sum_{I' I'_3} \omega_{I', I'_3} \int d^2 b dq_1 dq_2 \dots dq_n |\langle I' I'_3 n_+ n_- n_0 | \hat{S}(s, \vec{b}) | II_3 \rangle|^2, \quad (12)$$

$$n = n_+ + n_- + n_0$$

where N_{II_3} is the corresponding normalization factor determined by

$$\sum_{\{n\}} P_{II_3}(n_+, n_-, n_0) = 1.$$

This is our basic relation for calculating various pion-multiplicity distributions, pion multiplicities, and pion correlations between definite charge combinations. In general, the probability $P_{II_3}(n_+ n_- n_0)$ would depend on (I', I'_3) dynamically. The final-leading-particles tend to favor the $(I', I'_3) \approx (I, I_3)$ case. However, if the leading particles are colliding nuclei, an almost equal probability for various (I', I'_3) seems reasonable approximation owing to the large number of possible leading isobars in the final state. Then we can sum over all (I', I'_3) using the group theory alone.

Recent studies of heavy-ion collisions at the partonic level [26] argue that the central region is mainly dominated by gluon jets. The valence quarks of the incom-

ing particles which escape from the interaction region form the outgoing leading particle system. Since gluon's isospin is zero, it is very likely that total isospin of the produced pions in the central region is also zero. This picture is certainly true if the central region is free from valence quarks, the situation expected to appear at the extremely high collision energies.

Let us now assume that pions are produced both singly and through isovector clusters of the ρ type [24]. In this case, the most appropriate tool for studying various pion correlations is the generating function $G_{II_3}(z, n_-)$:

$$G_{II_3}(z, n_-) = \sum_{n_0, n_+} P_{II_3}(n_+, n_-, n_0) z^{n_0}, \quad (13)$$

from which we can calculate, for example

$$\langle n_0 \rangle_{n_-} = \frac{d}{dz} \ln G_{II_3}(1, n_-), \quad (14)$$

$$f_{2, n_-}^0 = \frac{d^2}{dz^2} \ln G_{II_3}(1, n_-), \quad (15)$$

and

$$P_{II_3}(n_0) = \frac{1}{n_0!} \frac{d^{n_0}}{dz^{n_0}} \sum_{n_-} G_{II_3}(0, n_-). \quad (16)$$

The form of this generating function is

$$G_{II_3}(z, n_-) = (I + \frac{1}{2}) \frac{(I - I_3)!}{(I + I_3)!} \int_{-1}^1 dx |P_I^{I_3}(x)|^2 \frac{A(z, x)^{n_0}}{n_0!} e^{-B(z, x)}, \quad (17)$$

where

$$2A(z, x) = (1 - x^2)\bar{n}_\pi + z(1 - x^2)\bar{n}_\rho + 2x^2\bar{n}_\rho \quad (18)$$

and

$$2B(z, x) = \bar{n}_\pi(1 + x^2 - 2zx^2) + \bar{n}_\rho(2 - z(1 - x^2)). \quad (19)$$

Here $\bar{n}_\pi$ denotes the average number of directly produced pions, and $\bar{n}_\rho$ denotes the average number of ρ -type clusters which decay into two short-range correlated

pions. The function $P_I^{I_3}(x)$ denotes the associate Legendre polinomial. Note that $A(1, x) = B(1, x)$.

The total number of emitted pions is

$$\bar{n} = \bar{n}_\pi + 2\bar{n}_\rho. \quad (20)$$

The behavior of $P(n_0) \equiv P_{00}(n_0)$ for $\bar{n} = 50$ and different combinations of $(\bar{n}_\pi, \bar{n}_\rho)$ is shown in Fig. 1. The behaviour of the n_0 -dispersion

$$D(n_0)_{n_-}^2 = f_{2,n_-} - \langle n_0 \rangle_{n_-} \quad (21)$$

for a given number of negative pions and different pairs of $(\bar{n}_\pi, \bar{n}_\rho)$ in the case of $I = I_3 = 1$ is shown in Fig. 2.. Note that f_{2,n_-} is a sensitive quaty of the pairing properties of the pions.

We see that DCC-type behavior is expected only for $\bar{n}_\pi \neq 0$ and $\bar{n}_\rho = 0$. that is in events without ρ -resonances. Recent estimate of the ratio of ρ - mesons to pions, at accelerator energies, is $\bar{n}_\rho = 0.10\bar{n}_\pi$ [6].

3. *DCC from a quantum nonlinear sigma model*

In this section we establish the relationship between the quantum nonlinear sigma model coupled to quarks and our coherent state eikonal model description of the pions emerging from the DCC.

As is well known, the Lagrangian for QCD with two light up and down quarks has an approximate global $SU(2)_L \times SU(2)_R$ symmetry, which at low temperatures, is spontaneously broken to $SU(2)_V$ by a nonzero value of the quark condensate $\langle \bar{q}_L q_R \rangle$, which is regarded as an order parameter of the system. This order parameter can be represented as a four-component vector $\phi \equiv (\sigma, \vec{\pi})$ buildt from the quark densities. The chiral symmetry then corresponds to $O(4)$ rotations in internal space.

The true vacuum of the theory is defined as $\langle\phi\rangle = (\langle\sigma\rangle, \vec{0})$, with $\langle\sigma\rangle \neq 0$. In QCD the spontaneous symmetry breakdown leads to nearly massless Goldstone bosons (the pions) and gives the constituent-quark mass. At low energies and large distances (momentum scale smaller than 1GeV) the dynamics of QCD is described by an effective Lagrangian containing the $\sigma, \vec{\pi}$ fields and constituent quarks.

In the DCC dynamics we distinguish three stages: formation, evolution and decay stage. In the conventional approach [19] one starts with a chirally symmetric phase at TT_c and DCC formation happens as T, due to a rapid expansion or cooling, drops below T_c spontaneously breaking the chiral symmetry.

The evolutionary stage of the DCC is usually described by the classical chiral dynamics based on the σ -model, mostly the linear σ - model. We consider the nonlinear σ -model coupled to quarks at zero temperature [35] which is expected to describe the late stage of the DCC evolution.

The Lagrangian for the nonlinear σ -model coupled to quarks is

$$L = \frac{f_\pi^2}{4} \text{Tr}(\partial_\mu U^\dagger \partial^\mu U) + \bar{q}(i\gamma\partial)q - g f_\pi \bar{q}Uq \quad (22)$$

where

$$U = \exp(i\gamma_5 \frac{\vec{\pi} \cdot \vec{\tau}}{f_\pi}) \quad (23)$$

We parametrize the pion field in the following form

$$\vec{\pi}(x) = f_\pi \vec{n}(x) \theta(x) \quad (24)$$

where $\vec{n}(x)$ is an unit vector which determines the isospin orientation of the pion field, obeying $\vec{n}^2 = 1$.

The Euler-Lagrange equations of motion for θ and $\vec{n}$ are

$$\square\theta - \sin\theta \cos\theta \partial_\mu \vec{n} \cdot \partial^\mu \vec{n} = -i \frac{m_Q}{f_\pi^2} \vec{n} \cdot (\bar{Q} \vec{\tau} \gamma_5 Q)$$

$$\partial_\mu(\sin^2\theta\vec{n} \times \partial^\mu\vec{n}) = -i\frac{m_Q}{f_\pi^2}\vec{n} \times (\bar{Q}\vec{\tau}\gamma_5 Q) \sin\theta \quad (25)$$

$$(26)$$

where Q denotes the constituent quark defined by

$$Q = e^{i\gamma_5 \frac{\vec{\pi} \cdot \vec{\tau}}{2f_\pi}} q \quad (27)$$

and $m_Q = gf_\pi$ is the constituent quark mass.

We treat

$$-i\frac{m_Q}{f_\pi}\bar{Q}\vec{\tau}\gamma_5 Q = \vec{j}(x) \quad (28)$$

as a given classical external source of pions and identify it with the source function $\vec{j}_1(s, \vec{b}; x)$ in our eikonal model. For the class of solutions that can be rotated into a uniform one, $\vec{n}(x) = \vec{n}$, known as the Anselm-class of solutions [16], the solutions with constant $\vec{n}$ can be realized if the source points to a certain fixed direction $\vec{n}$ in the isospace:

$$\vec{j}(x) = j(x)\vec{n} \quad (29)$$

Then the equation of motion for the pion field reduces to

$$\square\theta(x) = j(x) \quad (30)$$

with

$$\vec{\pi}(x) = f_\pi\theta(x)\vec{n} \quad (31)$$

This relationship offers us the possibility to study the importance of various quark sources in the DCC formation.

In quantum chiral field theory of the nonlinear σ -model the role of a strong coupling of $q\pi$ -interaction has the quantity m_Q/f_π . Since the Lagrangian of the nonlinear σ -model is nonpolynomial, a suitable renormalization procedure and operator

normal ordering should be formulated [29]. Let

$$\Phi =: e^{i\gamma_5 \frac{\vec{\pi}\vec{\tau}}{f_\pi}} - 1 : \quad (32)$$

denote the chiral super field of the pion. The Lagrangian describing the interaction of this super chiral field with quarks is now

$$L = \frac{f_\pi^2}{4} Tr(\partial_\mu \Phi \partial^\mu \Phi) + \bar{q}(i\gamma\partial - m_Q)q - \frac{m_Q}{f_\pi} \bar{q}\Phi q. \quad (33)$$

The chiral super propagator of the field Φ is defined by

$$\Delta_\Phi(x) = \langle T(\Phi(x)\Phi(0)) \rangle \quad (34)$$

Its explicite form, after number of algebraic manipulations, is

$$\Delta_\Phi(x) = \{1 \otimes 1 + \frac{f_\pi^2}{3}(\gamma_5 \vec{\tau}) \otimes (\gamma_5 \vec{\tau}) \partial_\Delta\} \partial_\Delta [\Delta ch(\frac{\Delta}{f_\pi^2})] + 1 \quad (35)$$

where

$$\langle T(\pi_i(x)\pi_j(0)) \rangle = \delta_{ij} \Delta(x). \quad (36)$$

and

$$\Delta(x) = \frac{1}{4\pi^2} \frac{1}{x^2 - i\epsilon}$$

.

The multipion exchange potential between two quarks is related to the Fourier transform of $\Delta_\Phi(x)$ in the following way:

$$\bar{u}(p'_1)\bar{u}(p'_2)\tilde{\Delta}_\Phi(q)u(p_1)u(p_2) = \omega'_1\omega'_2\tilde{V}_\Phi(q)\omega_1\omega_2 \quad (37)$$

where

$$u(p) = \sqrt{\frac{\epsilon_p + m}{\epsilon_p}} \begin{pmatrix} \omega \\ \omega' \end{pmatrix}$$

and

$$\omega' = \frac{\vec{\sigma}\vec{p}}{\epsilon_p + m}\omega$$

with $\omega^*\omega = 1$.

For studying the multipion production on quarks, it is necessary to find the correlator

$$\Delta_{nm}(q_1, q_2, \dots, q_n; l_1, l_2, \dots, l_m | x) = \langle n | T(\Phi(x)\Phi(0)) | m \rangle \quad (38)$$

This is an involved task and has not been done yet for arbitrary n and m .

In the case of soft chiral pion bremsstrahlung [37-40], in which every incoming and outgoing quark line is replaced by

$$q \longrightarrow \exp(i\gamma_5 \frac{\vec{\pi}\vec{\tau}}{2f_\pi})q \quad (39)$$

the distribution of neutral pions for large n is of the form $1/\sqrt{n_0 n}$ which is typical for coherent pion production without invoking the notion of DCC formation.

Therefore, we conclude that $nP(n_0 | n) \sim \sqrt{n/n_0}$ behaviour is not always a definite signature of DCC formation.

4. Conclusion

The results of the present analysis have shown the experimental observation of DCC is strongly affected by the ρ/π ratio. In particular, we have found that:

- Within the framework of an unitary eikonal model with factorization, energy conservation and global conservation of isospin the DCC-type fluctuation of the neutral pion fraction (f) could be observed if the ρ/π ratio is small in an event;

- The DCC effect also depends on isospin of the initial-leading- particle system;
- The coherent production of ρ -type clusters supresses the DCC-type behaviour;
- The factorization property of the scattering amplitude in the impact parameter space may be related to the isospin-uniform solutions of the quantum nonlinear σ -model coupled to quarks;
- The multipion exchange potential between two quarks can be found;
- The soft chiral pion bremsstrahlung also leads to anomalously large ratio of neutral to charged pions.

Acknowledgment This work was supported by the Ministry of Science of Croatia under Contract No. 0098004.

Figure captions :

Fig.1. The curves represent $P(n_0)$ for different combinations of $(\bar{n}_\pi, \bar{n}_\rho)$, the average number of singly produced pions and the average number of ρ -type clusters, respectively.

Fig. 2. The dispersion for two neutral pions as a function of the number of negative pions for $I = I_3 = 1$. The curves represent different combinations of $(\bar{n}_\pi, \bar{n}_\rho)$.

References

- [1] H. Heiselberg, G.A.Baym, B. Blättel, L.L. Frankfurt and M. Strikman, Phys. Rev. Lett. **67** (1991) 2946.
G. Baym, B. Blättel, L.L. Frankfurt, H. Heiselberg and M. Strikman, Phys. Rev. C **52** (1995) 1604.
- [2] G. Baym, G. Friedman and I. Sarcevic, Phys. Lett. B **219** (1989) 205.
H. Heiselberg, Phys. Rept. **351** (2001) 161.
- [3] M. Gazdzicki and S. Mrowczynski, Z. Phys C **54** (1992) 127.
- [4] K. Rajagopal and F. Wilczek, Nucl. Phys. B **399** (1993) 395.
S. Gavin, Nucl. Phys. A **590** (1995) 163c.
- [5] M. Stephanov, K. Rajagopal and E. Shuryak, Phys. Rev. Lett. **81** (1998) 4816;
Phys. Rev. D **60** (1999) 114028.
- [6] S. Jeon and V. Koch, Phys. Rev. Lett. **83** (1999) 5435.
S. Jeon and V. Koch, Phys. Rev. Lett. **85** (2000) 2076.
- [7] S. Mrowczynski, Phys. Lett. B **430** (1998) 9; Phys. Lett. B **459** (1999) 13.
S.A. Voloshin et al., Phys. Rev. C **60** (1999) 204901.

- [8] G. Baym and H. Heiselberg, Phys. Lett B **469** (1999) 7.
H. Heiselberg and A.D. Jackson, Phys. Rev. C **63** (2001) 064904.
- [9] C.M.G. Lattes, T. Fujimoto and S. Hasegawa, Phys. Rep. **65** (1980) 151.
- [10] D. Horn and R. Silver, Ann. Phys. **66** (1971) 509.
- [11] J.C. Botke, D.J. Scalapino and R.L. Sugar, Phys. Rev. D **9** (1974) 813.
- [12] P. Pirilä, Acta Phys. Pol. B **8** (1977) 305.
- [13] I.V. Andreev, Sov. Physica JETP Lett. **33** (1981) 367.
- [14] M. Martinis and V. Mikuta, Phys. Rev. D **12** (1975) 909.
- [15] L.J. Reinders, Acta Phys. Pol. B **7** (1976) 256.
- [16] A.A. Anselm, Phys. Lett. B **217** (1989) 169 .
A.A. Anselm and M. Ryskin, Phys. Lett. B **266** (1991) 482.
A.A. Anselm and M. Bander, JETP Lett. **59** (1994) 503.
A.A. Anselm and M. Ryskin, Z.Phys. A **358** (1997) 353.

- [17] J.D. Bjorken, Int. J. Mod. Phys. A **7** (1992) 4819; Acta Phys. Pol. B **23** (1992) 561.
J.D. Bjorken, K.L. Kowalski and C.C. Taylor, SLAC-PUB-6109 (April 1993); hep-ph/9309235.
- [18] J.P. Blaizot and A. Krzywicki, Phys. Rev. D **46** (1992) 246.
A. Krzywicki, Phys. Rev. D **48** (1993) 5190.
J.P. Blaizot and A. Krzywicki, Phys. Rev. D **50** (1994) 442.
J.P. Blaizot and A. Krzywicki, Acta Phys. Polon. B **27** (1996) 1687.
- [19] K.L. Kowalski and C.C. Taylor, hep-ph/9211282.
K. Rajagopal and F. Wilczek, Nucl. Phys. B **399** (1993) 395; Nucl. Phys. B **404** (1993) 577.
- [20] K. Rajagopal, hep-ph/9703258 (published in Hirschegg Workshop 1997, 136).
- [21] S. Pratt, Phys. Lett. B **301** (1993) 159.
- [22] G. Wilk and Z. Włodarczyk, Heavy Ion Phys. **4** (1996) 395.
- [23] MiniMax Collab., T.C. Brooks et al., Phys. Rev. D **61** (2000) 032003.
- [24] M. Martinis, V. Mikuta-Martinis, A. Švarc and J. Črnugelj, Fizika B **3** (1994) 197.

- M. Martinis, V. Mikuta-Martinis, A. Švarc and J. Črnugelj, Phys. Rev. D **51** (1995) 2482.
- M. Martinis, V. Mikuta-Martinis and J. Črnugelj, Phys. Rev. C **52** (1995) 1073.
- [25] V. Karmanov and A. Kudrjavev, ITEP-88, (1983).
- [26] R.D. Amado et al., Phys. Rev. Lett. **72** (1994) 970.
- [27] UA5 Collab., K.Alpgard et al., Phys. Lett. B **115** (1982) 71.
- [28] UA1 Collab., G. Arnison et al., Phys. Lett. B **122** (1983) 189.
- [29] UA5 Collab., G. Alner et al.: Phys. Lett. B **180** (1986) 415.
- [30] WA 98 Collab., M.M. Aggarwal et al., Phys. Lett. B **420** (1998) 169.
- [31] MiniMax Collab., T.C. Brooks et al., Phys. Rev. D **55** (1997) 5667.
- [32] B. Mohanty, T.K. Nayak, D.P. Mahapatra and Y.P. Viyogi, nucl-ex/0211007.
- [33] S. Gavin, Nucl. Phys. A **590** (1995) 163c.

- [34] M. Martinis, V. Mikuta-Martinis and J. Crnugelj, Acta Phys. Slov. **49** (1999) 875.
- [35] Z. Huang and M. Suzuki, Phys. Rev. D **53** (1996) 891.
M. Suzuki, Phys. Rev. D **54** (1996) 3556.
- [36] H. Leutwyler, Selected topics in nonperturbative QCD, Varenna 1995, 1 (hep-ph/9406283).
H. Leutwyler, Chiral dynamics workshop 1994, Cambridge, MA, 14 (hep-ph/9409423).
- [37] S. Weinberg, Phys. Rev. D **2** (1970) 674.
- [38] L.S. Brown, Phys. Rev. D **2** (1970) 3083.
- [39] N. Bilić and M. Martinis, Fizika **6** (1974) 145.
- [40] I.V. Andreev et al., Phys. Lett. B **388** (1996) 197.



